



If you are taking Rifampin, you should know that:

- ♦ your urine, saliva, or tears can become orange in color.
- ♦ your soft contact lenses may get stained. This can happen ***very quickly!*** Do not wear soft contact lenses when taking rifampin.
- ♦ your skin may become more sensitive to the sun. Use sunscreen and cover skin that might be exposed to the sun.
- ♦ some birth control methods **do not work as well** when you are taking rifampin. Women should use a back-up form of birth control, such as condoms, while taking rifampin.



Protect yourself, your family and friends from TB. Finish your TB treatment!

If you have questions, please contact:

---

---

---

---



<http://www.vdh.state.va.us/epi/tb>



November 8, 2002

# What You Should Know About






# Taking Tuberculosis Medicines


## Taking medicines to kill TB germs can cure TB disease.


TB germs are strong. You need to take medicine for many months to be sure all of the TB germs are dead.





Remember the following when taking your TB pills:

-  Follow your health care provider's directions for taking your pills.
-  Remember to take your pills. Your health care worker will talk to you about **Directly Observed Therapy (DOT)**. DOT is when a health care worker is with you when you take your TB pills. DOT can be done at a time and place that works for you. DOT makes it easy to take your TB pills.
-  Do not stop taking your pills too soon. Even though you might start to feel better in a few weeks, TB can come back. You can again spread TB to others.

 Tell your health care worker if you are taking any other medicines. This includes medicine, vitamins and herbs that you buy without a prescription.

 Avoid drinking beer, wine or liquor while taking TB medicines.

 Tell your health care provider if you are pregnant, breastfeeding or taking birth control pills.

 Tell your health care worker if you have any health problems other than TB.

---

✓ The TB medicine(s) you are taking:

\_\_\_\_\_ Isoniazid (INH)

\_\_\_\_\_ Rifampin

\_\_\_\_\_ Pyrazinamide (PZA)

\_\_\_\_\_ Ethambutol

\_\_\_\_\_ Other medicine \_\_\_\_\_

\_\_\_\_\_ Other medicine \_\_\_\_\_

\_\_\_\_\_ Other medicine \_\_\_\_\_

**TB medicines are usually safe.** Very few people have side effects. If you have any of the problems listed below, you must call or see your health care worker right away.

- ▶ Loss of appetite
- ▶ Vomiting
- ▶ Stomach pain
- ▶ Nausea
- ▶ Yellowing of your eyes or skin
- ▶ Tingling in your fingers or toes
- ▶ Tingling or numbness around your mouth
- ▶ Blurred vision or change in your vision
- ▶ Dizziness
- ▶ Aching joints
- ▶ Fever for more than 3 days
- ▶ Skin rash
- ▶ Bleeding easily
- ▶ Bruising easily
- ▶ \_\_\_\_\_
- ▶ \_\_\_\_\_
- ▶ \_\_\_\_\_
- ▶ \_\_\_\_\_
- ▶ \_\_\_\_\_
- ▶ \_\_\_\_\_